

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A01N 43/04, A61K 31/70, C12N 15/00, 15/63	A1	(11) International Publication Number: WO 00/38518 (43) International Publication Date: 6 July 2000 (06.07.00)
(21) International Application Number: PCT/US99/31093 (22) International Filing Date: 28 December 1999 (28.12.99) (30) Priority Data: 60/113,923 28 December 1998 (28.12.98) US (71) Applicant: ARCH DEVELOPMENT CORPORATION [US/US]; The University of Chicago, Suite 405, 5640 South Ellis, Chicago, IL 60637 (US). (72) Inventors: LEIDEN, Jeffrey, M.; Harvard Medical School, HSPH Building II, Room 117, 677 Huntington Avenue, Boston, MA 02115 (US). SVENSSON, Eric; University of Chicago, Dept. of Medicine, 5841 South Maryland, Chicago, IL 60637 (US). (74) Agents: BRAINARD, Charles, R. et al.; Kenyon & Kenyon, One Broadway, New York, NY 10004 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: EFFICIENT AND STABLE <i>IN VIVO</i> GENE TRANSFER TO CARDIOMYOCYTES USING RECOMBINANT ADENO-ASSOCIATED VIRUS VECTORS (57) Abstract This invention relates to the use of recombinant adeno-associated virus (rAAV) vectors to transduce cardiomyocytes <i>in vivo</i> by infusing the rAAV into a coronary artery or coronary sinus. rAAV infection is not associated with detectable myocardial inflammation or myocyte necrosis. Thus, rAAV is a useful vector for the stable expression of therapeutic genes in the myocardium and can be used to deliver genes for inducing angiogenesis, inhibiting angiogenesis, stimulating cell proliferation, inhibiting cell proliferation and/or treating or ameliorating other cardiovascular conditions.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

EFFICIENT AND STABLE *IN VIVO* GENE TRANSFER TO CARDIOMYOCYTES USING RECOMBINANT ADENO-ASSOCIATED VIRUS VECTORS

5 This application is a continuation-in-part application of provisional application
U.S. Ser. No. 60/113,923, filed December 28, 1998, which is incorporated by reference
herein in its entirety.

This work was supported in part by grants from the National Institutes of Health
(DK-48997, AR-42895, and HL-54592) to Jeffrey M. Leiden.

10 FIELD OF THE INVENTION

The ability to stably and efficiently program recombinant gene expression in
cardiomyocytes facilitates gene therapy approaches for a variety of cardiovascular
diseases and conditions. Accordingly, this invention relates to the use of recombinant
adeno-associated virus (rAAV) vectors to transduce cardiomyocytes *in vivo* by infusing
15 the rAAV into a coronary artery or coronary sinus. For example, coronary artery
perfusion of mouse hearts with a rAAV encoding the LacZ gene produced efficient
transduction of cardiomyocytes which was stable for at least 8 weeks. Moreover, rAAV
infection is not associated with detectable myocardial inflammation or myocyte necrosis.
Thus, rAAV is a useful vector for the stable expression of therapeutic genes in the
20 myocardium and can be used to deliver genes for inducing angiogenesis, inhibiting
angiogenesis, stimulating cell proliferation, inhibiting cell proliferation and/or treating or
ameliorating other cardiovascular conditions.

BACKGROUND OF THE INVENTION

25 Myocardial gene therapy can be used for the treatment of a number of
cardiovascular diseases, including ischemic cardiomyopathies, congestive heart failure,
and malignant arrhythmias (Nabel (1995) *Circulation* 91:541-548). A useful vector for
myocardial gene delivery will allow efficient and stable transduction of cardiomyocytes
with a variety of transgenes after either direct intramyocardial injection or infusion into
30 the coronary arteries or sinuses. For example, plasmid DNA vectors injected directly into
the left ventricular myocardium have been expressed for ≥ 6 months by cardiomyocytes

adjacent to the area of injection (Lin et al. (1990a) *Circulation* 82:2217-2221; Kass et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:11498-11502; and Guzman et al. (1993) *Circ. Res.* 73:1202-1207). However, the therapeutic usefulness of this approach has been limited by the low efficiency of cardiomyocyte transduction (0.1% to 1.0% of cardiomyocytes in the area of injection).

Both intramyocardial injection and intracoronary infusion of replication-defective adenovirus (RDAd) vectors have been used to efficiently transduce cardiomyocytes in rodents, rabbits, and pigs *in vivo*. However, the feasibility of adenovirus-mediated gene transfer has been limited by immune responses to viral and foreign transgene proteins, which cause significant myocardial inflammation, eliminate virus-transduced cells within 30 days of infection, and thereby result in transient recombinant gene expression in immunocompetent hosts (Guzman et al. (1993) *Circ. Res.* 73:1202-1207; French et al. (1994) *Circulation* 90:2414-2424; and Barr et al. (1994) *Gene Ther.* 1:51-58).

Recently, rAAV vectors have been shown to program efficient and stable recombinant gene expression in skeletal muscle and liver in both rodents and primates (Fisher et al. (1997) *Nat. Med.* 3:306-312; Kessler et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:14082-14087; and Snyder et al. (1997) *Nat. Genet.* 16: 270-276) and in cardiac muscle directly injected with rAAV (U.S. Patent No. 5,858,351 to Podsakoff et al.). However, since rAAV vectors used in gene therapy applications, unlike RDAd, do not encode viral proteins, the rAAV vectors have not been associated with immune responses to foreign transgene proteins.

While a previous report showed that rAAV can transduce cardiomyocytes *in vivo*, the efficiency of rAAV-mediated transgene expression in the heart was both low (about 0.2%) and localized (Kaplitt et al. (1996) *Ann. Thorac. Surg.* 62:1669-1676). In that study, pigs hearts were rapidly perfused with a low titer of rAAV (less than 10^4 expressing units AAV per gram of body weight). Based on those results, infusing rAAV into the heart would have severely limited use as a vector for myocardial gene therapy. However, as demonstrated herein, this invention establishes that by infusing rAAV in much higher amounts proportional to body weight of the animal and for particular time periods, then rAAV provides unexpected efficient and stable gene transfer into the heart, opening up use of rAAV vectors to deliver therapeutically-effective molecules to

cardiomyocytes in amounts useful for treating or ameliorating cardiac diseases or conditions.

SUMMARY OF THE INVENTION

5 The present invention is directed to a method of treating a cardiovascular condition by infusing an rAAV vector into a coronary artery or a coronary sinus for a time and in an amount sufficient to stably and efficiently transduce the cardiomyocytes perfused by the artery or sinus. The rAAV vector encodes at least one nucleic acid which is operably linked to a control region and which encodes a therapeutically-effective
10 molecule. After infusion and transduction of the cardiomyocytes, the therapeutically-effective molecule is expressed in an amount effective to treat or ameliorate the cardiovascular condition.

 Thus, this method provides a means of delivering AAV vectors in a stable and efficient manner. The vector can be infused by any convenient means and in conjunction
15 with surgery or other cardiac procedure, if desired.

BRIEF DESCRIPTION OF THE DRAWINGS

 Figure 1. Schematic of AAV_{CMV-LacZ}. ITR indicates inverted terminal repeats; BGH pA, bovine growth hormone polyadenylation signal; CMV Pr, CMV immediate-
20 early promoter; *LacZ*, bacterial *LacZ* gene.

 Figure 2. Gene transfer into cardiomyocytes in vivo with AAV_{CMV-LacZ}. Gross sections (left) and photomicrographs (right) of mouse hearts after coronary artery perfusion with 1.5×10^9 IU of AAV_{CMV-LacZ} and staining with X-gal. Bar=25 microns.

DETAILED DESCRIPTION OF THE INVENTION

25 This invention relates to treating cardiovascular conditions using rAAV vectors. In accordance with the invention, an rAAV vector encoding a therapeutically effective molecule is infused into a coronary artery or a coronary sinus to deliver the vector to the heart in a manner which stably and efficiently transduces cardiomyocytes. It has
30 unexpectedly been found that the ability to obtain stable and efficient transduction of

cardiomyocytes by rAAV depends upon the duration of the infusion period and the amount of virus infused relative to body weight.

Moreover, rAAV displays significant advantages for myocardial gene transfer compared with plasmid DNA or adenovirus vectors. For example, rAAV, when delivered as described herein, allows efficient transduction of cardiomyocytes. Further, rAAV vectors program stable expression of foreign transgenes in immunocompetent hosts. The stability of transgene expression observed with rAAV even after expression of a foreign transgene protein likely reflects the fact that rAAV vectors, unlike their adenovirus counterparts, do not express any viral gene products and are therefore significantly less immunogenic. This lack of immunogenicity represents a major advantage of rAAV for myocardial gene transfer.

Hence, the invention is directed to a method of treating a cardiovascular condition which comprises infusing an rAAV vector into a coronary artery or sinus of an animal for a time and in an amount sufficient to stably and efficiently transduce cardiomyocytes perfused by the artery or sinus, wherein that vector encodes at least one nucleic acid, *i.e.*, the transgene, encoding a therapeutically-effective molecule; and expressing the therapeutically-effective molecule in an amount effective to treat or ameliorate the cardiovascular condition. Further, the nucleic acid is operably linked to a control region, *e.g.*, promoters, enhancers, termination signals and the like, to permit expression of the molecule. When more than one nucleic acid is present on the rAAV vector, each can be controlled separately by individual control regions or, any group of them, or all of them, can be controlled in an operon, *i.e.*, with one control region driving expression of multiple genes on a single transcript.

rAAV vectors useful in the present invention can be any rAAV vector with one or more transgenes (or nucleic acids of interest) inserted therein in a manner allowing expression of the transgene under control of appropriate regulatory elements such as promoters, enhancers, transcription terminators and the like. rAAV vectors are well known in the art and can be prepared by standard methodology known to those of ordinary skill in the art. For example, U.S. Patent No. 5,858,351 and the references cited therein describe a variety of rAAV vectors suitable for use in gene therapy as well as how to make and propagate those vectors (*see, e.g.*, Kotin (1994) Human Gene Therapy

5:793-801 or Berns, "Parvoviridae and their Replication" in Fundamental Virology, 2nd Edition, (Fields & Knipe, eds.)).

A "transgene" or "nucleic acid of interest" or the "nucleic acid encoded in the rAAV vector" as used herein refers to any nucleotide sequence which encodes a therapeutically-effective molecule that can be used to treat a cardiovascular condition. Such transgenes may normally be foreign to the animal being treated or may be a gene normally found in that animal for which altered expression (e.g., temporal, spatial or amount of expression) is desired to achieve a particular therapeutic effect. The therapeutically-effective molecule encoded by the transgene is protein or an anti-sense RNA that imparts a benefit to the animal or subject undergoing treatment or amelioration of a cardiac condition or disease in accordance with this invention.

Proteins that can be administered to treat or ameliorate cardiovascular conditions are numerous and include, but are not limited to, molecules competent to induce angiogenesis, e.g., angiogenesis factors; anti-angiogenesis factors; proteins capable of inhibiting vascular smooth muscle cell proliferation; proteins useful for treating atherosclerosis; proteins useful for treating restenosis, proteins useful for stimulating cardiomyocyte activity; proteins capable of secretion from cardiomyocytes that exert their effect in the heart or capable of transport to other locales for treatment of a cardiovascular condition or disease; hormones, cytokines or growth factors useful for treating cardiac conditions or diseases; and proteins capable of stimulating vascular smooth muscle cell proliferation. Other genes encoding proteins useful in this invention include ion channel genes, contractile protein genes, phospholamban encoding genes and genes encoding β adrenergic receptors or β adrenergic kinases.

Angiogenic factors include, but are not limited to FGF-1, FGF-2, FGF-5, VEGF, HIF-1 and the like. Proteins useful for treating restenosis include thymidine kinase, cytosine deaminase, p21, p27, p53, Rb, and NF- κ B. Hence, this invention can be used to deliver any protein via an rAAV vector that has a therapeutic benefit for treating or ameliorating a cardiovascular condition or disease.

A protein competent to induce angiogenesis or an "angiogenesis factor" as used herein is a protein or substance that causes proliferation of new blood vessels and includes fibroblast growth factors, endothelial cell growth factors or other proteins with such

biological activity. Particular proteins known to induce angiogenesis are FGF-1, FGF-2, FGF-5, VEGF and active fragment thereof, and HIF-1. Proteins competent to inhibit angiogenesis or "anti-angiogenesis factors" are proteins or substances that inhibit the formation of new blood vessels.

5 Anti-sense RNA that can be administered to treat or ameliorate cardiovascular conditions have one of the same activities as proteins useful in the invention. Such RNA include, but are not limited to, c-myc, c-myc and others. Anti-sense RNA molecules, including how to design and use such molecules in expression vectors are well known in the art and can be constructed by routine methodology. Thus a strand of RNA whose
10 sequence of bases is complementary to the sense, or translated, RNA strand can form a duplex to block translation or degradation of a particular mRNA or otherwise control or alter expression of the desired mRNA.

 As used herein, a "control region" or "regulatory element" refers to polyadenylation signals, upstream regulatory domains, promoters, enhancers, transcription
15 termination sequences and the like which regulate the transcription and translation of a nucleic acid sequence.

 The term "operably linked" refers to an arrangement of elements wherein the components are arranged so as to perform their usual function. Thus, control regions or regulatory elements operably linked to a coding sequence are capable of effecting the
20 expression of the coding sequence. The control elements need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

25 The regulatory elements of the invention can be derived from any source, e.g., viruses, mammals, insects or even synthetic, provided that they function in cardiomyocytes. For example, any promoter can be used to control expression of the transgene. Such promoters can be promiscuous, i.e., active in many cell types, such as the SV40 early promoter, the mouse mammary tumor virus LTR promoter, the adenovirus
30 major late promoter (Ad MLP), a herpes simplex promoter, a CMV promoter such as the

CMV immediate early promoter, a rous sarcoma virus (RSV) promoter. Alternatively the promoter can be tissue-specific for expression in cardiomyocytes.

The rAAV is delivered to cardiac myocytes by infusion into a coronary artery or coronary sinus. This mode of delivery has also been referred to as intraluminal delivery through a coronary artery, intracoronary delivery or intraarterial delivery. As used herein, infusion into a coronary artery includes intracoronary perfusion. In accordance with the invention, the rAAV vector can be infused when the heart is in situ, i.e., in the body cavity or when the heart or heart tissue (cardiac tissue) has been removed from the body as might occur when the heart is being donated for transplant into a recipient. In the case of a heart being prepared for transplantation or for heart tissue, the vector can be infused through any artery or vein attached thereto, by contacting with or soaking the heart in an appropriately concentrated solution of the vector, or by a combination of both. Thus as described herein, infusion includes delivering rAAV to a heart or heart tissue ex vivo by the means disclosed herein. If necessary, the infusion can be repeated at intervals such as 3 months, 6 months, one year, or as appropriately determined.

As used herein, treating cardiac conditions include treating cardiac or cardiovascular diseases. Examples of cardiac conditions subject to treatment or amelioration according to the method of the present invention include, but are not limited to, myocardial ischemia, myocardial infarction, congestive heart failure, dilated and hypertrophic cardiomyopathy, cardiac arrhythmia, cardiac hypertrophy, cardiac transplantation and rejection. For example, if the cardiac condition, such as ischemia, can be treated or improved by inducing angiogenesis, then the rAAV vector used in accordance with the method of this invention would encode an angiogenesis factor.

Thus the rAAV vector is infused into a coronary artery for a time and in an amount sufficient to stably and efficiently transduce cardiac tissue perfused by the artery, wherein the AAV vector encodes a therapeutically-effective molecule which is expressed in the cardiac tissue in an amount effective to treat or ameliorate a cardiovascular condition including, but not limited to, inducing angiogenesis, inhibiting angiogenesis, stimulating or inhibiting cell proliferation, treating restenosis, treating atherosclerosis, treating congestive heart failure, treating ischemic cardiomyopathies or treating malignant

arrhythmias, myocardial infarction, congestive heart failure, or dilated and hypertrophic cardiomyopathy.

The method of the present invention can be used with any animal, including but not limited to, mammals such as rodents, dogs, cats, cattle, primates and humans.

5 Preferably the method is used for gene therapy to treat human acquired or inherited cardiac conditions or diseases.

The present invention thus provides a method of treating and/or ameliorating a cardiovascular condition by infusing an rAAV vector for a time and in an amount sufficient to stably and efficiently transduce cardiomyocytes which was heretofore
10 unachievable by methods known in the art. For this invention, stable and efficient transduction means that a significant number of cardiomyocytes are transduced and are capable of expressing the protein for a prolonged period of time. Stable and efficient transduction occurs over a period of time and can actually be observed over time as an increase in the percentage of transduced cardiomyocytes, as continued expression of the
15 transgene, or as continued observation of the therapeutic effect at a molecular, microscopic or macroscopic level. For example, with angiogenesis, stable and efficient transduction can be manifested by ongoing development and or growth of new blood vessels, by observing the improved blood flow to the heart, or by determining measuring the level of ischemia in the heart tissue.

20 Alternatively, efficient transduction occurs when at least about 10%, and preferably more, of the cardiomyocytes have been transduced, i.e., infected by, the rAAV. By following the methods of the invention and by observing at particular times after transduction ranging over a few to many weeks, about 25%, about 40% or even about 50% of the cardiomyocytes will be transduced. While about 10% of the cardiomyocytes
25 can be transduced using only rAAV, this percentage can be increased by co-infusing adenovirus as a helper virus without adverse effects.

The time of infusion contributes to achieving stable and efficient transduction of the cardiomyocytes as well. Thus the infusion time ranges from about 2 minutes to about 30 minutes, more preferably from about 5 minutes to about 20 minutes and most
30 preferably is about fifteen minutes.

The amount of rAAV infused into the animal is proportional to the body weight of the animal. Hence in accordance with the invention, stable and efficient transduction occurs when the amount of rAAV infused ranges from about 1×10^5 IU (infectious units) of AAV per gram body weight to about 1×10^9 IU AAV per gram body weight, and preferably from about 1×10^6 IU AAV per gram body weight to about 1×10^8 IU AAV per gram body weight, and most preferably is about $5-6 \times 10^7$ IU AAV per gram body weight.

The example described below demonstrates efficient and stable transduction of cardiac myocytes *in vivo* after intracoronary infusion of an rAAV vector.

Throughout this application, various publications, patents, and patent applications have been referred to. The teachings and disclosures of these publications, patents, and patent applications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which the present invention pertains.

It is to be understood and expected that variations in the principles of invention herein disclosed in an exemplary embodiment may be made by one skilled in the art and it is intended that such modifications, changes, and substitutions are to be included within the scope of the present invention.

EXAMPLE

Intracoronary Infusion of rAAV

I. Methods:

5 Plasmids and Viruses

The structure of pAAV_{CMV-LacZ} is shown in Figure 1. Ad_{CMV-LacZ} and the E3-deleted adenovirus, Ad_{d1309}, were propagated and purified as described (Barr 1994).

Propagation and Purification of rAAV

10 rAAV was prepared as described (Rolling et al. (1995) *Mol. Biotechnol.* 3:9-15) and purified by cesium chloride gradient centrifugation. Viral titer was assessed by a dot blot hybridization assay to determine the number of viral genomes per milliliter and by infecting HeLa cells with the virus and staining with X-gal 24 hours after infection. All viral preparations had titers of 1 to 2×10^{11} genomes/mL, and 2 to 3×10^9 infectious units (IU)/mL.

15 Intracoronary Perfusion With rAAV

Adult C57BL/6 mouse hearts were perfused via the left carotid artery with cardioplegia solution (110 mmol/L NaCl, 25 mmol/L KCl, 22 mmol/L NaHCO₃, 16 mmol/L MgCl₂, 0.8 mmol/L CaCl₂, 40 mmol/L glucose) at 4°C until they stopped beating. They were then perfused ex vivo for 15 minutes with 1.5×10^9 IU of AAV_{CMV-LacZ} in 0.5 mL of PBS at a rate of 33 μ L/min at 4°C. After perfusion, the hearts were transplanted into the neck of a syngeneic host with anastomosis of the donor aorta to the right common carotid artery of the host and anastomosis of the donor pulmonary artery to the right external jugular vein (Lin et al. (1990b) *J. Heart Transplant.* 9:720-723) (n=3 for each time point).

25 X-Gal Staining

Freshly isolated hearts were fixed in PBS plus 1.25% glutaraldehyde for 10 minutes at room temperature, stained overnight with X-gal (Lin 1990a), and counterstained with eosin.

30

β -Galactosidase Activity

Cardiac homogenates were assayed for β -galactosidase (β -gal) activity and protein concentrations. β -Gal activities were normalized for total protein and for the number of infectious rAAV or RDAd particles injected.

5 **II. Results**

Many clinical applications of myocardial gene therapy may require the stable and efficient transduction of cardiomyocytes distributed throughout large areas of myocardium. Coronary artery infusions of RDAd have been shown to result in the efficient transduction of cardiomyocytes throughout the region of perfused myocardium
10 (Barr 1994). To test whether rAAV is similarly capable of transducing cardiomyocytes after coronary artery perfusion, hearts from C57BL/6 mice were explanted and perfused with 1.5×10^9 IU of AAV_{CMV-LacZ} for 15 minutes at 4°C via a catheter placed in the left common carotid artery. These perfused hearts were then transplanted into syngeneic hosts, and the arterial circulation was reestablished by anastomosis of the transplanted
15 aorta to the recipient carotid artery. Such transplanted and revascularized hearts resumed beating and continued to do so until the recipient mice were killed 2, 4, or 8 weeks after perfusion. Two weeks after perfusion, small numbers (<1 %) of β -gal-positive cardiomyocytes were detected throughout the myocardium of the rAAV-perfused hearts (Figure 2). By 4 weeks after perfusion, $\approx 40\%$ of the cardiomyocytes were β -gal positive.
20 This high level of transduction was stable at weeks after perfusion, with >50% of the cardiomyocytes continuing to express β -gal. Similar increases in recombinant gene expression over the first several weeks after rAAV infection have been observed in skeletal muscle (Fisher 1997; Kessler 1998). It has been postulated that such increases may reflect the gradual process of conversion of the single-stranded AAV genome into a
25 double-stranded DNA molecule that is competent for transcription of the transgene (Ferrari et al. (1996) *J. Virol.* 70:3227-3234). Thus, rAAV delivered by coronary artery perfusion can be used to stably transduce cardiomyocytes throughout the myocardium.

WE CLAIM:

1. A method of treating a cardiovascular condition which comprises:
infusing a recombinant adeno-associated virus (AAV) vector into a coronary
artery or a coronary sinus for a time and in an amount sufficient to stably and efficiently
transduce cardiomyocytes perfused by said artery or said sinus, wherein said AAV vector
5 encodes at least one nucleic acid operably linked to a control region, said nucleic acid
encoding a therapeutically-effective molecule; and
expressing said therapeutically-effective molecule in an amount effective to treat
or ameliorate said cardiovascular condition.
- 10 2. The method of Claim 1, wherein said AAV transduces at least about 10 % of
said cardiomyocytes.
3. The method of Claim 1, wherein said AAV transduces at least about 40% of
said cardiomyocytes.
4. The method of Claim 1, wherein said AAV transduces at least about 50% of
15 said cardiomyocytes.
5. The method of Claim 1, wherein said AAV is infused for at least about 2
minutes to about 30 minutes.
6. The method of Claim 1, wherein said AAV is infused for at least about 5
minutes to about 20 minutes.
- 20 7. The method of Claim 1, wherein said AAV is infused for about 15 minutes.
8. The method of Claim 1, wherein said amount of AAV is about 1×10^5 IU AAV
per gram body weight to about 1×10^9 IU AAV per gram body weight.
9. The method of Claim 9, wherein said amount of AAV is about 1×10^6 IU AAV
per gram body weight to about 1×10^8 IU AAV per gram body weight.
- 25 10. The method of Claim 9, wherein said amount of AAV is about 6×10^7 IU
AAV per gram body weight.
11. The method of Claim 1, wherein about 1×10^5 IU AAV per gram body weight
to about 1×10^9 IU AAV per gram body weight is infused for about 2 to about 30 minutes.
12. The method of Claim 11, wherein about 1×10^6 IU AAV per gram body
30 weight to about 1×10^8 IU AAV per gram body weight is infused.

13. The method of Claim 11, wherein about 6×10^7 IU AAV per gram body weight is infused.

14. The method of any one of Claims 11, 12 or 13, wherein said AAV is infused for about 5 to about 20 minutes.

5 15. The method of any one of Claims 11, 12 or 13, wherein said AAV is infused for about 15 minutes.

16. The method of Claim 11, wherein about 6×10^7 IU AAV per gram body weight is infused for about 15 minutes.

10 17. The method of Claim 1, wherein said coronary artery is infused *ex vivo or in vivo*.

18. The method of Claim 1, wherein said therapeutically-effective molecule is an anti-sense RNA or a protein.


15 19. The method of Claim 1 wherein therapeutically-effective molecule is an ion channel gene, a contractile protein, a phospholamban, a β adrenergic receptor, a β adrenergic kinase, a growth factor, an angiogenic factor, a protein or nucleic acid capable of inducing angiogenesis, or a protein or nucleic acid capable of inhibiting angiogenesis.

20. The method of Claim 1, wherein said therapeutically-effective molecule is FGF-1, FGF-2, FGF-5, VEGF, or HIF-1.


20 21. The method of Claim 1, wherein said therapeutically-effective molecule is thymidine kinase, p21, p27, p53, Rb or NF- κ B.

22. The method of Claim 1, wherein said cardiovascular condition is restenosis, atherosclerosis, congestive heart failure, ischemic cardiomyopathy, malignant arrhythmia, myocardial infarction, congestive heart failure, or dilated and hypertrophic cardiomyopathy.


25 23. The method of Claim 1, wherein treating or ameliorating said cardiovascular condition is for inducing angiogenesis, inhibiting angiogenesis, stimulating or inhibiting cell proliferation, treating restenosis, treating atherosclerosis, treating congestive heart failure, treating ischemic cardiomyopathy or treating malignant arrhythmia.



2 weeks



4 weeks



8 weeks



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/31093

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A01N 43/04; A61K 31/70; C12N 15/00, 15/63

US CL : 514/44; 435/320.1, 455, 456, 69.1, 325

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/44; 435/320.1, 455, 456, 69.1, 325

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	KAPLITT et al. Long-Term Gene Transfer in Porcine Myocardium After Coronary Infusion of an Adeno-Associated Virus Vector. Annals of Thoracic Surgery. December 1996, Vol. 62, No. 6, pages 1669-1676, see entire document.	1-23
Y	GNATENKO et al. Characterization of Recombinant Adeno-Associated Virus-2 as a Vehicle for Gene Delivery and Expression into Vascular Cells. Journal of Investigative Medicine. February 1997, Vol. 45, No. 2, pages 87-98, see entire document.	1-23

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

10 MARCH 2000

Date of mailing of the international search report

10 APR 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20540Authorized officer
JILL D. MARTINJOYCE BRIDGERS
PARALEGAL SPECIALIST
CHEMICAL PATENT

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/31093

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	SVENSSON et al. Efficient and Stable Transduction of Cardiomyocytes After Intramyocardial Injection of Intracoronary Perfusion With Recombinant Adeno-Associated Virus Vectors. Circulation. 19 January 1999, Vol. 99, No. 2, pages 201-205, see entire document.	1-23
Y,P	ALEXANDER et al. Gene Transfer and Models of Gene Therapy For The Myocardium. Clinical and Experimental Pharmacology and Physiology. September 1999, Vol. 26, No. 9, pages 661-668, see entire document.	1-23

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/31093

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

NPL, WEST, MEDLINE, CAPLUS, EMBASE, BIOSIS

search terms: adeno-associated virus, AAV, coronary artery, vessel, lumen, angiogenesis, cardiomyocytes, coronary sinus, vector, cardiovascular, ischemia, myocardium

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 July 2000 (06.07.2000)

PCT

(10) International Publication Number
WO 00/38518 A1

(51) International Patent Classification⁷: **A01N 43/04.**
A61K 31/70, C12N 15/00, 15/63

(21) International Application Number: PCT/US99/31093

(22) International Filing Date:
28 December 1999 (28.12.1999)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/113,923 28 December 1998 (28.12.1998) US

(71) Applicant: **ARCH DEVELOPMENT CORPORATION** [US/US]; The University of Chicago, Suite 405, 5640 South Ellis, Chicago, IL 60637 (US).

(72) Inventors: **LEIDEN, Jeffrey, M.**; Harvard Medical School, HSPH Building II, Room 117, 677 Huntington Avenue, Boston, MA 02115 (US). **SVENSSON, Eric**; University of Chicago, Dept. of Medicine, 5841 South Maryland, Chicago, IL 60637 (US).

(74) Agent: **WILSON, M., Lisa**; Hale and Dorr, LLP, 300 Park Avenue, New York, NY 10022 (US).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(48) Date of publication of this corrected version:
11 April 2002

(15) Information about Correction:
see PCT Gazette No. 15/2002 of 11 April 2002, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: EFFICIENT AND STABLE (*IN VIVO*) GENE TRANSFER TO CARDIOMYOCYTES USING RECOMBINANT ADENO-ASSOCIATED VIRUS VECTORS

(57) Abstract: This invention relates to the use of recombinant adeno-associated virus (rAAV) vectors to transduce cardiomyocytes *in vivo* by infusing the rAAV into a coronary artery or coronary sinus. rAAV infection is not associated with detectable myocardial inflammation or myocyte necrosis. Thus, rAAV is a useful vector for the stable expression of therapeutic genes in the myocardium and can be used to deliver genes for inducing angiogenesis, inhibiting angiogenesis, stimulating cell proliferation, inhibiting cell

WO 00/38518 A1

EFFICIENT AND STABLE *IN VIVO* GENE TRANSFER TO CARDIOMYOCYTES
USING RECOMBINANT ADENO-ASSOCIATED VIRUS VECTORS

5 This application is a continuation-in-part application of provisional application
U.S. Ser. No. 60/113,923, filed December 28, 1998, which is incorporated by reference
herein in its entirety.

This work was supported in part by grants from the National Institutes of Health
(DK-48997, AR-42895, and HL-54592) to Jeffrey M. Leiden.

10 FIELD OF THE INVENTION

The ability to stably and efficiently program recombinant gene expression in
cardiomyocytes facilitates gene therapy approaches for a variety of cardiovascular
diseases and conditions. Accordingly, this invention relates to the use of recombinant
adeno-associated virus (rAAV) vectors to transduce cardiomyocytes *in vivo* by infusing
15 the rAAV into a coronary artery or coronary sinus. For example, coronary artery
perfusion of mouse hearts with a rAAV encoding the LacZ gene produced efficient
transduction of cardiomyocytes which was stable for at least 8 weeks. Moreover, rAAV
infection is not associated with detectable myocardial inflammation or myocyte necrosis.
Thus, rAAV is a useful vector for the stable expression of therapeutic genes in the
20 myocardium and can be used to deliver genes for inducing angiogenesis, inhibiting
angiogenesis, stimulating cell proliferation, inhibiting cell proliferation and/or treating or
ameliorating other cardiovascular conditions.

BACKGROUND OF THE INVENTION

25 Myocardial gene therapy can be used for the treatment of a number of
cardiovascular diseases, including ischemic cardiomyopathies, congestive heart failure,
and malignant arrhythmias (Nabel (1995) *Circulation* 91:541-548). A useful vector for
myocardial gene delivery will allow efficient and stable transduction of cardiomyocytes
with a variety of transgenes after either direct intramyocardial injection or infusion into
30 the coronary arteries or sinuses. For example, plasmid DNA vectors injected directly into
the left ventricular myocardium have been expressed for ≥ 6 months by cardiomyocytes

adjacent to the area of injection (Lin et al. (1990a) *Circulation* 82:2217-2221; Kass et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:11498-11502; and Guzman et al. (1993) *Circ. Res.* 73:1202-1207). However, the therapeutic usefulness of this approach has been limited by the low efficiency of cardiomyocyte transduction (0.1% to 1.0% of cardiomyocytes in the area of injection).

Both intramyocardial injection and intracoronary infusion of replication-defective adenovirus (RDAd) vectors have been used to efficiently transduce cardiomyocytes in rodents, rabbits, and pigs *in vivo*. However, the feasibility of adenovirus-mediated gene transfer has been limited by immune responses to viral and foreign transgene proteins, which cause significant myocardial inflammation, eliminate virus-transduced cells within 30 days of infection, and thereby result in transient recombinant gene expression in immunocompetent hosts (Guzman et al. (1993) *Circ. Res.* 73:1202-1207; French et al. (1994) *Circulation* 90:2414-2424; and Barr et al. (1994) *Gene Ther.* 1:51-58).

Recently, rAAV vectors have been shown to program efficient and stable recombinant gene expression in skeletal muscle and liver in both rodents and primates (Fisher et al. (1997) *Nat. Med.* 3:306-312; Kessler et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:14082-14087; and Snyder et al. (1997) *Nat. Genet.* 16: 270-276) and in cardiac muscle directly injected with rAAV (U.S. Patent No. 5,858,351 to Podsakoff et al.). However, since rAAV vectors used in gene therapy applications, unlike RDAd, do not encode viral proteins, the rAAV vectors have not been associated with immune responses to foreign transgene proteins.

While a previous report showed that rAAV can transduce cardiomyocytes *in vivo*, the efficiency of rAAV-mediated transgene expression in the heart was both low (about 0.2%) and localized (Kaplitt et al. (1996) *Ann. Thorac. Surg.* 62:1669-1676). In that study, pigs hearts were rapidly perfused with a low titer of rAAV (less than 10^4 expressing units AAV per gram of body weight). Based on those results, infusing rAAV into the heart would have severely limited use as a vector for myocardial gene therapy. However, as demonstrated herein, this invention establishes that by infusing rAAV in much higher amounts proportional to body weight of the animal and for particular time periods, then rAAV provides unexpected efficient and stable gene transfer into the heart, opening up use of rAAV vectors to deliver therapeutically-effective molecules to

cardiomyocytes in amounts useful for treating or ameliorating cardiac diseases or conditions.

SUMMARY OF THE INVENTION

5 The present invention is directed to a method of treating a cardiovascular condition by infusing an rAAV vector into a coronary artery or a coronary sinus for a time and in an amount sufficient to stably and efficiently transduce the cardiomyocytes perfused by the artery or sinus. The rAAV vector encodes at least one nucleic acid which is operably linked to a control region and which encodes a therapeutically-effective
10 molecule. After infusion and transduction of the cardiomyocytes, the therapeutically-effective molecule is expressed in an amount effective to treat or ameliorate the cardiovascular condition.

 Thus, this method provides a means of delivering AAV vectors in a stable and efficient manner. The vector can be infused by any convenient means and in conjunction
15 with surgery or other cardiac procedure, if desired.

BRIEF DESCRIPTION OF THE DRAWINGS

 Figure 1. Schematic of AAV_{CMV-LacZ}. ITR indicates inverted terminal repeats; BGH pA, bovine growth hormone polyadenylation signal; CMV Pr, CMV immediate-
20 early promoter; *LacZ*, bacterial *LacZ* gene.

 Figure 2. Gene transfer into cardiomyocytes in vivo with AAV_{CMV-LacZ}. Gross sections (left) and photomicrographs (right) of mouse hearts after coronary artery perfusion with 1.5×10^9 IU of AAV_{CMV-LacZ} and staining with X-gal. Bar=25 microns.

DETAILED DESCRIPTION OF THE INVENTION

25 This invention relates to treating cardiovascular conditions using rAAV vectors. In accordance with the invention, an rAAV vector encoding a therapeutically effective molecule is infused into a coronary artery or a coronary sinus to deliver the vector to the heart in a manner which stably and efficiently transduces cardiomyocytes. It has
30 unexpectedly been found that the ability to obtain stable and efficient transduction of

cardiomyocytes by rAAV depends upon the duration of the infusion period and the amount of virus infused relative to body weight.

Moreover, rAAV displays significant advantages for myocardial gene transfer compared with plasmid DNA or adenovirus vectors. For example, rAAV, when delivered as described herein, allows efficient transduction of cardiomyocytes. Further, rAAV vectors program stable expression of foreign transgenes in immunocompetent hosts. The stability of transgene expression observed with rAAV even after expression of a foreign transgene protein likely reflects the fact that rAAV vectors, unlike their adenovirus counterparts, do not express any viral gene products and are therefore significantly less immunogenic. This lack of immunogenicity represents a major advantage of rAAV for myocardial gene transfer.

Hence, the invention is directed to a method of treating a cardiovascular condition which comprises infusing an rAAV vector into a coronary artery or sinus of an animal for a time and in an amount sufficient to stably and efficiently transduce cardiomyocytes perfused by the artery or sinus, wherein that vector encodes at least one nucleic acid, *i.e.*, the transgene, encoding a therapeutically-effective molecule; and expressing the therapeutically-effective molecule in an amount effective to treat or ameliorate the cardiovascular condition. Further, the nucleic acid is operably linked to a control region, *e.g.*, promoters, enhancers, termination signals and the like, to permit expression of the molecule. When more than one nucleic acid is present on the rAAV vector, each can be controlled separately by individual control regions or, any group of them, or all of them, can be controlled in an operon, *i.e.*, with one control region driving expression of multiple genes on a single transcript.

rAAV vectors useful in the present invention can be any rAAV vector with one or more transgenes (or nucleic acids of interest) inserted therein in a manner allowing expression of the transgene under control of appropriate regulatory elements such as promoters, enhancers, transcription terminators and the like. rAAV vectors are well known in the art and can be prepared by standard methodology known to those of ordinary skill in the art. For example, U.S. Patent No. 5,858,351 and the references cited therein describe a variety of rAAV vectors suitable for use in gene therapy as well as how to make and propagate those vectors (*see, e.g.*, Kotin (1994) Human Gene Therapy

5:793-801 or Berns, "Parvoviridae and their Replication" in Fundamental Virology, 2nd Edition, (Fields & Knipe, eds.)).

A "transgene" or "nucleic acid of interest" or the "nucleic acid encoded in the rAAV vector" as used herein refers to any nucleotide sequence which encodes a therapeutically-effective molecule that can be used to treat a cardiovascular condition. Such transgenes may normally be foreign to the animal being treated or may be a gene normally found in that animal for which altered expression (e.g., temporal, spatial or amount of expression) is desired to achieve a particular therapeutic effect. The therapeutically-effective molecule encoded by the transgene is protein or an anti-sense RNA that imparts a benefit to the animal or subject undergoing treatment or amelioration of a cardiac condition or disease in accordance with this invention.

Proteins that can be administered to treat or ameliorate cardiovascular conditions are numerous and include, but are not limited to, molecules competent to induce angiogenesis, e.g., angiogenesis factors; anti-angiogenesis factors; proteins capable of inhibiting vascular smooth muscle cell proliferation; proteins useful for treating atherosclerosis; proteins useful for treating restenosis, proteins useful for stimulating cardiomyocyte activity; proteins capable of secretion from cardiomyocytes that exert their effect in the heart or capable of transport to other locales for treatment of a cardiovascular condition or disease; hormones, cytokines or growth factors useful for treating cardiac conditions or diseases; and proteins capable of stimulating vascular smooth muscle cell proliferation. Other genes encoding proteins useful in this invention include ion channel genes, contractile protein genes, phospholamban encoding genes and genes encoding β adrenergic receptors or β adrenergic kinases.

Angiogenic factors include, but are not limited to FGF-1, FGF-2, FGF-5, VEGF, HIF-1 and the like. Proteins useful for treating restenosis include thymidine kinase, cytosine deaminase, p21, p27, p53, Rb, and NF- κ B. Hence, this invention can be used to deliver any protein via an rAAV vector that has a therapeutic benefit for treating or ameliorating a cardiovascular condition or disease.

A protein competent to induce angiogenesis or an "angiogenesis factor" as used herein is a protein or substance that causes proliferation of new blood vessels and includes fibroblast growth factors, endothelial cell growth factors or other proteins with such

biological activity. Particular proteins known to induce angiogenesis are FGF-1, FGF-2, FGF-5, VEGF and active fragment thereof, and HIF-1. Proteins competent to inhibit angiogenesis or "anti-angiogenesis factors" are proteins or substances that inhibit the formation of new blood vessels.

5 Anti-sense RNA that can be administered to treat or ameliorate cardiovascular conditions have one of the same activities as proteins useful in the invention. Such RNA include, but are not limited to, c-myb, c-myc and others. Anti-sense RNA molecules, including how to design and use such molecules in expression vectors are well know in the art and can be constructed by routine methodology. Thus a strand of RNA whose
10 sequence of bases is complementary to the sense, or translated, RNA strand can form a duplex to block translation or degradation of a particular mRNA or otherwise control or alter expression of the desired mRNA.

 As used herein, a "control region" or "regulatory element" refers to polyadenylation signals, upstream regulatory domains, promoters, enhancers, transcription
15 termination sequences and the like which regulate the transcription and translation of a nucleic acid sequence.

 The term "operably linked" refers to an arrangement of elements wherein the components are arranged so as to perform their usual function. Thus, control regions or regulatory elements operably linked to a coding sequence are capable of effecting the
20 expression of the coding sequence. The control elements need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

25 The regulatory elements of the invention can be derived from any source, e.g., viruses, mammals, insects or even synthetic, provided that they function in cardiomyocytes. For example, any promoter can used to control expression of the transgene. Such promoters can be promiscuous, i.e., active in many cell types, such as the SV40 early promoter, the mouse mammary tumor virus LTR promoter, the adenovirus
30 major late promoter (Ad MLP), a herpes simplex promoter, a CMV promoter such as the

CMV immediate early promoter, a rous sarcoma virus (RSV) promoter. Alternatively the promoter can be tissue-specific for expression in cardiomyocytes.

5 The rAAV is delivered to cardiac myocytes by infusion into a coronary artery or coronary sinus. This mode of delivery has also been referred to as intraluminal delivery through a coronary artery, intracoronary delivery or intraarterial delivery. As used herein, infusion into a coronary artery includes intracoronary perfusion. In accordance with the invention, the rAAV vector can be infused when the heart is in situ, i.e., in the body cavity or when the heart or heart tissue (cardiac tissue) has been removed from the body as might occur when the heart is being donated for transplant into a recipient. In the case
10 of a heart being prepared for transplantation or for heart tissue, the vector can be infused through any artery or vein attached thereto, by contacting with or soaking the heart in an appropriately concentrated solution of the vector, or by a combination of both. Thus as described herein, infusion includes delivering rAAV to a heart or heart tissue ex vivo by the means disclosed herein. If necessary, the infusion can be repeated at intervals such as
15 3 months, 6 months, one year, or as appropriately determined.

As used herein, treating cardiac conditions include treating cardiac or cardiovascular diseases. Examples of cardiac conditions subject to treatment or amelioration according to the method of the present invention include, but are not limited to, myocardial ischemia, myocardial infarction, congestive heart failure, dilated and
20 hypertrophic cardiomyopathy, cardiac arrhythmia, cardiac hypertrophy, cardiac transplantation and rejection. For example, if the cardiac condition, such as ischemia, can be treated or improved by inducing angiogenesis, then the rAAV vector used in accordance with the method of this invention would encode an angiogenesis factor.

Thus the rAAV vector is infused into a coronary artery for a time and in an
25 amount sufficient to stably and efficiently transduce cardiac tissue perfused by the artery, wherein the AAV vector encodes a therapeutically-effective molecule which is expressed in the cardiac tissue in an amount effective to treat or ameliorate a cardiovascular condition including, but not limited to, inducing angiogenesis, inhibiting angiogenesis, stimulating or inhibiting cell proliferation, treating restenosis, treating atherosclerosis,
30 treating congestive heart failure, treating ischemic cardiomyopathies or treating malignant

arrhythmias, myocardial infarction, congestive heart failure, or dilated and hypertrophic cardiomyopathy.

The method of the present invention can be used with any animal, including but not limited to, mammals such as rodents, dogs, cats, cattle, primates and humans.

5 Preferably the method is used for gene therapy to treat human acquired or inherited cardiac conditions or diseases.

The present invention thus provides a method of treating and/or ameliorating a cardiovascular condition by infusing an rAAV vector for a time and in an amount sufficient to stably and efficiently transduce cardiomyocytes which was heretofore
10 unachievable by methods known in the art. For this invention, stable and efficient transduction means that a significant number of cardiomyocytes are transduced and are capable of expressing the protein for a prolonged period of time. Stable and efficient transduction occurs over a period of time and can actually be observed over time as an increase in the percentage of transduced cardiomyocytes, as continued expression of the
15 transgene, or as continued observation of the therapeutic effect at a molecular, microscopic or macroscopic level. For example, with angiogenesis, stable and efficient transduction can be manifested by ongoing development and or growth of new blood vessels, by observing the improved blood flow to the heart, or by determining measuring the level of ischemia in the heart tissue.

20 Alternatively, efficient transduction occurs when at least about 10%, and preferably more, of the cardiomyocytes have been transduced, i.e., infected by, the rAAV. By following the methods of the invention and by observing at particular times after transduction ranging over a few to many weeks, about 25%, about 40% or even about 50% of the cardiomyocytes will be transduced. While about 10% of the cardiomyocytes
25 can be transduced using only rAAV, this percentage can be increased by co-infusing adenovirus as a helper virus without adverse effects.

The time of infusion contributes to achieving stable and efficient transduction of the cardiomyocytes as well. Thus the infusion time ranges from about 2 minutes to about 30 minutes, more preferably from about 5 minutes to about 20 minutes and most
30 preferably is about fifteen minutes.

The amount of rAAV infused into the animal is proportional to the body weight of the animal. Hence in accordance with the invention, stable and efficient transduction occurs when the amount of rAAV infused ranges from about 1×10^5 IU (infectious units) of AAV per gram body weight to about 1×10^9 IU AAV per gram body weight, and preferably from about 1×10^6 IU AAV per gram body weight to about 1×10^8 IU AAV per gram body weight, and most preferably is about $5-6 \times 10^7$ IU AAV per gram body weight.

The example described below demonstrates efficient and stable transduction of cardiac myocytes *in vivo* after intracoronary infusion of an rAAV vector.

Throughout this application, various publications, patents, and patent applications have been referred to. The teachings and disclosures of these publications, patents, and patent applications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which the present invention pertains.

It is to be understood and expected that variations in the principles of invention herein disclosed in an exemplary embodiment may be made by one skilled in the art and it is intended that such modifications, changes, and substitutions are to be included within the scope of the present invention.

EXAMPLE

Intracoronary Infusion of rAAV

I. Methods:

5 Plasmids and Viruses

The structure of pAAV_{CMV-LacZ} is shown in Figure 1. Ad_{CMV-LacZ} and the E3-deleted adenovirus, Ad_{d1309}, were propagated and purified as described (Barr 1994).

Propagation and Purification of rAAV

10 rAAV was prepared as described (Rolling et al. (1995) *Mol. Biotechnol.* 3:9-15) and purified by cesium chloride gradient centrifugation. Viral titer was assessed by a dot blot hybridization assay to determine the number of viral genomes per milliliter and by infecting HeLa cells with the virus and staining with X-gal 24 hours after infection. All viral preparations had titers of 1 to 2×10^{11} genomes/mL, and 2 to 3×10^9 infectious units (IU)/mL.

15 Intracoronary Perfusion With rAAV

Adult C57BL/6 mouse hearts were perfused via the left carotid artery with cardioplegia solution (110 mmol/L NaCl, 25 mmol/L KCl, 22 mmol/L NaHCO₃, 16 mmol/L MgCl₂, 0.8 mmol/L CaCl₂, 40 mmol/L glucose) at 4°C until they stopped beating. They were then perfused ex vivo for 15 minutes with 1.5×10^9 IU of AAV_{CMV-LacZ} in 0.5 mL of PBS at a rate of 33 μ L/min at 4°C. After perfusion, the hearts were transplanted into the neck of a syngeneic host with anastomosis of the donor aorta to the right common carotid artery of the host and anastomosis of the donor pulmonary artery to the right external jugular vein (Lin et al. (1990b) *J. Heart Transplant.* 9:720-723) (n=3 for each time point).

25 X-Gal Staining

Freshly isolated hearts were fixed in PBS plus 1.25% glutaraldehyde for 10 minutes at room temperature, stained overnight with X-gal (Lin 1990a), and counterstained with eosin.

30

β -Galactosidase Activity

Cardiac homogenates were assayed for β -galactosidase (β -gal) activity and protein concentrations. β -Gal activities were normalized for total protein and for the number of infectious rAAV or RDAd particles injected.

5 **II. Results**

Many clinical applications of myocardial gene therapy may require the stable and efficient transduction of cardiomyocytes distributed throughout large areas of myocardium. Coronary artery infusions of RDAd have been shown to result in the efficient transduction of cardiomyocytes throughout the region of perfused myocardium (Barr 1994). To test whether rAAV is similarly capable of transducing cardiomyocytes after coronary artery perfusion, hearts from C57BL/6 mice were explanted and perfused with 1.5×10^9 IU of AAV_{CMV-LacZ} for 15 minutes at 4°C via a catheter placed in the left common carotid artery. These perfused hearts were then transplanted into syngeneic hosts, and the arterial circulation was reestablished by anastomosis of the transplanted aorta to the recipient carotid artery. Such transplanted and revascularized hearts resumed beating and continued to do so until the recipient mice were killed 2, 4, or 8 weeks after perfusion. Two weeks after perfusion, small numbers (<1 %) of β -gal-positive cardiomyocytes were detected throughout the myocardium of the rAAV-perfused hearts (Figure 2). By 4 weeks after perfusion, \approx 40% of the cardiomyocytes were β -gal positive. This high level of transduction was stable at weeks after perfusion, with >50% of the cardiomyocytes continuing to express β -gal. Similar increases in recombinant gene expression over the first several weeks after rAAV infection have been observed in skeletal muscle (Fisher 1997; Kessler 1998). It has been postulated that such increases may reflect the gradual process of conversion of the single-stranded AAV genome into a double-stranded DNA molecule that is competent for transcription of the transgene (Ferrari et al. (1996) *J. Virol.* 70:3227-3234). Thus, rAAV delivered by coronary artery perfusion can be used to stably transduce cardiomyocytes throughout the myocardium.

WE CLAIM:

1. A method of treating a cardiovascular condition which comprises:
infusing a recombinant adeno-associated virus (AAV) vector into a coronary artery or a coronary sinus for a time and in an amount sufficient to stably and efficiently transduce cardiomyocytes perfused by said artery or said sinus, wherein said AAV vector encodes at least one nucleic acid operably linked to a control region, said nucleic acid encoding a therapeutically-effective molecule; and
expressing said therapeutically-effective molecule in an amount effective to treat or ameliorate said cardiovascular condition.
2. The method of Claim 1, wherein said AAV transduces at least about 10 % of said cardiomyocytes.
3. The method of Claim 1, wherein said AAV transduces at least about 40% of said cardiomyocytes.
4. The method of Claim 1, wherein said AAV transduces at least about 50% of said cardiomyocytes.
5. The method of Claim 1, wherein said AAV is infused for at least about 2 minutes to about 30 minutes.
6. The method of Claim 1, wherein said AAV is infused for at least about 5 minutes to about 20 minutes.
7. The method of Claim 1, wherein said AAV is infused for about 15 minutes.
8. The method of Claim 1, wherein said amount of AAV is about 1×10^5 IU AAV per gram body weight to about 1×10^9 IU AAV per gram body weight.
9. The method of Claim 9, wherein said amount of AAV is about 1×10^6 IU AAV per gram body weight to about 1×10^8 IU AAV per gram body weight.
10. The method of Claim 9, wherein said amount of AAV is about 6×10^7 IU AAV per gram body weight.
11. The method of Claim 1, wherein about 1×10^5 IU AAV per gram body weight to about 1×10^9 IU AAV per gram body weight is infused for about 2 to about 30 minutes.
12. The method of Claim 11, wherein about 1×10^6 IU AAV per gram body weight to about 1×10^8 IU AAV per gram body weight is infused.

13. The method of Claim 11, wherein about 6×10^7 IU AAV per gram body weight is infused.

14. The method of any one of Claims 11, 12 or 13, wherein said AAV is infused for about 5 to about 20 minutes.

5 15. The method of any one of Claims 11, 12 or 13, wherein said AAV is infused for about 15 minutes.

16. The method of Claim 11, wherein about 6×10^7 IU AAV per gram body weight is infused for about 15 minutes.

10 17. The method of Claim 1, wherein said coronary artery is infused *ex vivo* or *in vivo*.

18. The method of Claim 1, wherein said therapeutically-effective molecule is an anti-sense RNA or a protein.

15 19. The method of Claim 1 wherein therapeutically-effective molecule is an ion channel gene, a contractile protein, a phospholamban, a β adrenergic receptor, a β adrenergic kinase, a growth factor, an angiogenic factor, a protein or nucleic acid capable of inducing angiogenesis, or a protein or nucleic acid capable of inhibiting angiogenesis.


20. The method of Claim 1, wherein said therapeutically-effective molecule is FGF-1, FGF-2, FGF-5, VEGF, or HIF-1.

20 21. The method of Claim 1, wherein said therapeutically-effective molecule is thymidine kinase, p21, p27, p53, Rb or NF- κ B.

22. The method of Claim 1, wherein said cardiovascular condition is restenosis, atherosclerosis, congestive heart failure, ischemic cardiomyopathy, malignant arrhythmia, myocardial infarction, congestive heart failure, or dilated and hypertrophic cardiomyopathy.

25 23. The method of Claim 1, wherein treating or ameliorating said cardiovascular condition is for inducing angiogenesis, inhibiting angiogenesis, stimulating or inhibiting cell proliferation, treating restenosis, treating atherosclerosis, treating congestive heart failure, treating ischemic cardiomyopathy or treating malignant arrhythmia.

FIG. 1



2/2

FIG. 2



2 weeks

4 weeks

8 weeks

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/31093

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A01N 43/04; A61K 31/70; C12N 15/00, 15/63

US CL : 514/44; 435/320.1, 455, 456, 69.1, 325

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/44; 435/320.1, 455, 456, 69.1, 325

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	KAPLITT et al. Long-Term Gene Transfer in Porcine Myocardium After Coronary Infusion of an Adeno-Associated Virus Vector. Annals of Thoracic Surgery. December 1996, Vol. 62, No. 6, pages 1669-1676, see entire document.	1-23
Y	GNATENKO et al. Characterization of Recombinant Adeno-Associated Virus-2 as a Vehicle for Gene Delivery and Expression into Vascular Cells. Journal of Investigative Medicine. February 1997, Vol. 45, No. 2, pages 87-98, see entire document.	1-23

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

10 MARCH 2000

Date of mailing of the international search report

10 APR 2000

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

 Authorized officer
 JILL D. MARTIN

 JOYCE BRIDGERS
 PARALEGAL SPECIALIST
 CHEMICAL MATRIX ✓

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/31093

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	SVENSSON et al. Efficient and Stable Transduction of Cardiomyocytes After Intramyocardial Injection of Intracoronary Perfusion With Recombinant Adeno-Associated Virus Vectors. Circulation. 19 January 1999, Vol. 99, No. 2, pages 201-205, see entire document.	1-23
Y,P	ALEXANDER et al. Gene Transfer and Models of Gene Therapy For The Myocardium. Clinical and Experimental Pharmacology and Physiology. September 1999, Vol. 26, No. 9, pages 661-668, see entire document.	1-23

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/31093

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

NPL, WEST, MEDLINE, CAPLUS, EMBASE, BIOSIS

search terms: adeno-associated virus, AAV, coronary artery, vessel, lumen, angiogenesis, cardiomyocytes, coronary sinus, vector, cardiovascular, ischemia, myocardium

